ELSEVIER

Fluid Phase Equilibria 136 (1997) 377–378

FLUID PHASE EQUILIBRIA

Contents of Volume 136 (1997)

Vol. 136, Nos. 1–2

Preface vii

On the derivation of thermodynamic equilibrium criteria
J.L. De Medeiros and M.J. Estillac de Mello Cardoso (Rio de Janeiro, Brazil) 1–13
Modeling of thermodynamically difficult systems
S.I. Sandler, J.P. Wolbach, M. Castier and G. Escobedo-Alvarado (Newark, DE, USA) 15–29
Electrostatic model for substituent and solvent effects on organic acids
X.-G. Hu, H.-X. Zong and R.-S. Lin (Hangzhou, China) 31–36
Pressure and temperature dependence of excess thermodynamic quantities of Lennard–Jones binary mixtures
F. Kawaizumi, M. Ohba, T. Fukuyama and H. Nomura (Nagoya-shi, Japan) 37–47
A refined lattice–hole theory with group contribution for unified description of single- and multi-component real fluids
K.-P. Yoo, S.J. Yoo and C.S. Lee (Seoul, Korea) 49–62
Compilation of group-contribution prediction of VLE and excess enthalpy
K. Kojima (Tokyo, Japan) 63–77
Calculation of combinatory entropy in complex polymer solutions based on the Flory–Huggins theory
S. Saeki (Fukui, Japan) 79–85
Calculation of phase diagram of polymer solution over supercritical fluid region based on the Patterson-Delmas theory of corresponding state
S. Saeki (Fukui, Japan) 87–92
Phase separations for mixtures in well-characterized porous materials: Liquid–liquid transitions
M. Sliwinska-Bartkowiak, R. Sikorski, S.L. Sowers, L.D. Gelb and K.E. Gubbins (Poznan, Poland, Ithaca, NY, USA) . . 93–109
On the unique behavior of molar excess enthalpies for the supercritical fluid mixtures
I. Fujihara, Y. Shimada and K. Nakanishi (Osaka, Japan, Okayama) 111–114
Simulation of non-equilibrium physical adsorption taking into account the adatoms interactions
K.T. Kholmurodov, V.K. Fedyanin and I.V. Puzynin (Moscow, Russia) 115–133
Measurement and correlation of infinite dilution activity coefficients of ethers in alkanes
K. Fukuchi, K. Miyoshi and Y. Arai (Ube, Japan, Fukuoka) 135–139
Liquid–liquid equilibria of ternary systems containing alkane, methanol, and ether
H. Higashiuchi, T. Watanabe and Y. Arai (Kurume, Japan, Omuta, Fukuoka) 141–146
VLE modelling for aqueous systems containing methyldiethanolamine, carbon dioxide and hydrogen sulfide
G. Kuranov, B. Rumpf, G. Maurer and N. Smirnova (St. Petersburg, Russian Federation, Kaiserslautern, Germany) 147–162
Swelling equilibria of poly(*N*-isopropylacrylamide) gel in glucose and starch aqueous solution
T. Ishidao, I.-S. Song, N. Ohtani, K. Sato, Y. Iwai and Y. Arai (Fukuoka, Japan) 163–171
Micellar solution–solid phase equilibrium for ternary systems containing two *N*-alkanoyl-*N*-methylglucamine homologues
M. Alexeeva, T. Churjusova, N. Smirnova and A. Vlasov (St. Petersburg, Russia) 173–183
A new fibre optic-based technique for the spectrophotometric measurement of phase and chemical equilibria in aqueous solutions of volatile weak electrolytes
M.A. Siddiqi, J. Krissmann and K. Lucas (Duisburg, Germany) 185–195
The enthalpic stabilization on molecular inclusion of butanediol isomers into cyclodextrin cavities
M. Fujisawa, T. Kimura and S. Takagi (Wakayama, Japan, Osaka) 197–205
'Intermolecular interactions' in aqueous solutions of three components including lysozyme
Y. Koga, P. Westh, C. Trandum and C.A. Haynes (Vancouver, BC, Canada) 207–221

Thermodynamic properties of polyoxyethyleneglycol dimethyl ether + *n*-alkane mixtures
C.A. Tovar, E. Carballo, C.A. Cerdeiriña, M.I. Paz Andrade and L. Romaní (Orense, Spain, Santiago de Compostela) . . . 223–234
Thermodynamic properties of the mixture of methoxyethanol and cyclohexane: Measurements at the temperatures 293.15, 298.15 and 303.15 K above and below UCST
M. Nishimoto, S. Tabata, K. Tamura and S. Murakami (Osaka, Japan) . . . 235–247
Excess enthalpies of ethyl acetate and butyl acetate with 2-alkoxyethanols at 298.15 K
D. Venkatesulu, P. Venkatesu and M.V. Prabhakara Rao (Andhra Pradesh, India) . . . 249–255
Excess enthalpies of binary mixtures {$x$dioxane isomer + $(1-x)$non-polar liquid} at 298.15 K
T. Takigawa, H. Ogawa, K. Tamura and S. Murakami (Osaka, Japan, Saitama) . . . 257–267
Excess enthalpies of *o*- and *m*-dihalogenated benzene + methyl methyl thiomethyl sulfoxide, + dimethyl sulfoxide at 298.15 K
T. Kimura, K. Suzuki and S. Takagi (Osaka, Japan) . . . 269–278
Thermodynamic properties of rigid polycyclic molecules. 1: Enthalpies of solution of fused ring polycyclic aromatic hydrocarbons
H. Ogawa, S. Murakami, T. Takigawa and M. Ohba (Saitama, Japan, Osaka, Nagoya-shi) . . . 279–287
Thermodynamic properties of rigid polycyclic molecules. (2) Partial molar volumes of polycyclic aromatics compared with the RISM integral equation theory
M. Ohba, T. Takigawa, H. Ogawa, S. Murakami and H. Nomura (Nagoya-shi, Japan, Osaka, Saitama) . . . 289–297
Excess molar volumes of (3-methyl-2-oxazolidinone + an alkan-1-ol) at 298.15 K, 313.15 K, and 328.15 K
Y. Uosaki, T. Hamaguchi and T. Moriyoshi (Tokushima, Japan) . . . 299–306
Excess properties of binary systems formed by carbon tetrachloride with isopropanol and n-butanol
L. Simoiu, I. Trandafir, M. Pleniceanu and M. Baniceru (Craiova, Romania) . . . 307–314
Excess molar volumes of ternary mixtures of di-*n*-butylether + 1-heptanol + *n*-octane at the temperature of 298.15 K
J.L. Legido, E. Jiménez, C. Franjo, L. Segade and M.I. Paz Andrade (Coruña, Spain, Vigo, Santiago de Compostela) . . . 315–321
Excess volumes of some halogenated aromatics + methyl methylthiomethyl sulfoxide, + dimethyl sulfoxide at 298.15 K
T. Kimura, Y. Sugihara and S. Takagi (Osaka, Japan) . . . 323–331
Adsorption from ternary liquid mixtures on silica gel of different mesoporosity
J. Goworek, A. Nieradka and A. Dąbrowski (Lublin, Poland) . . . 333–343
Combination of overall reaction rate with Gibbs energy minimization
P. Koukkari, I. Laukkanen and S. Liukkonen (Espoo, Finland) . . . 345–362
Predicting fluid viscosity by entropy production
Z. Suoqi (Beijing, China) . . . 363–371

Subject Index . . . 373–374

Author Index . . . 375